

## How weight and BMI affect getting pregnant



### What BMI means in fertility care

BMI is calculated from weight and height and is commonly used to classify adults as underweight, healthy weight, overweight, or obese. In many clinical settings, a BMI below 18.5 kg/m<sup>2</sup> is considered underweight, 18.5 to 24.9 is considered in the healthy range, 25 to 29.9 is considered overweight, and 30 or above is considered obesity. These categories are broad screening tools, not precise measurements of metabolic health, body composition, or reproductive potential.

In fertility care, BMI may be considered because body fat is metabolically active. Adipose tissue produces hormones and inflammatory mediators and influences insulin sensitivity, estrogen metabolism, and the hypothalamic-pituitary-gonadal axis, the hormonal system that regulates . However, BMI does not distinguish muscle from fat, does not capture fat distribution, and may be less accurate across ethnic groups, athletic body types, and people with certain medical conditions.

A medically literate way to think about BMI is this: it can identify patterns of risk at the population level, but predict with certainty whether one individual will or will not conceive. If your BMI is outside the typical

reference range, it is reasonable to discuss it as one part of a broader fertility assessment rather than as a standalone explanation.

### **Low weight, hormonal signaling, and ovulation**

Being underweight can reduce fertility by altering the hormonal signals required for regular . When the body has insufficient energy availability, the brain may reduce pulsatile gonadotropin-releasing hormone signaling. This can lower luteinizing hormone and follicle-stimulating hormone activity, leading to irregular cycles, anovulation, or absent periods.

The Better Health Channel notes that underweight women with a BMI below 18.5 may experience hormone imbalances that disrupt ovulation and can take longer than women in a healthy weight range to conceive. The American Society for Reproductive Medicine also states that people with low BMI may have irregular menstrual cycles and may stop ovulating.

Low weight may be related to many factors, including high physical training load, gastrointestinal disease, eating disorders, chronic stress, restrictive eating patterns, endocrine conditions, or other illnesses. Because the causes can be complex, it is safer to seek individualized care rather than attempting rapid weight gain without guidance. A clinician may consider menstrual history, nutrition, exercise, thyroid function, prolactin, signs of hypothalamic amenorrhea, and other reproductive hormone markers.

### **Higher BMI, ovulatory function, and time to pregnancy**

Overweight and obesity can also affect , particularly through ovulatory dysfunction. Excess adipose tissue can contribute to insulin resistance, elevated androgen levels, chronic low-grade inflammation, and decreased estrogen production. These factors may interfere with follicle development and . In some people, higher BMI coexists with polycystic ovary syndrome, or PCOS, although not everyone with a higher BMI has PCOS and not everyone with PCOS has a higher BMI.

A study using data from the United States National Health and Nutrition Examination Survey 2013-2018 found a non-linear association between BMI and female infertility. Below a BMI of 19.5 kg/m<sup>2</sup>, each unit increase in BMI was associated with a lower infertility risk; at or above 19.5 kg/m<sup>2</sup>, each unit

increase was associated with a small increase in infertility risk. After adjustment for other variables, the reported odds ratio was 1.03 per BMI unit. The same study noted interactions with factors such as age, marital status, diabetes history, and previous pregnancy, underscoring that BMI operates within a broader health context.

From a practical standpoint, higher BMI may be associated with longer time to pregnancy, more irregular cycles, and reduced likelihood of spontaneous in some individuals. But this is not universal. Many people with overweight or obesity without assistance. If cycles are consistently irregular, very long, absent, or unpredictable, it is worth discussing evaluation with a healthcare professional.

### **BMI and assisted reproductive treatment**

Weight outcomes with treatment, including induction and in vitro fertilization, or IVF. According to the American Society for Reproductive Medicine, overweight and obesity are associated with reduced natural rates and lower IVF success rates, including lower pregnancy rates in some treatment populations. Higher BMI may also affect medication dosing, ovarian response, egg retrieval logistics, anesthesia risk, and embryo transfer planning.

For people with low BMI, treatment may also be affected if  $\text{GnRH}$  is suppressed by inadequate energy availability or if overall is compromised. A clinic may recommend nutritional optimization before treatment, particularly if periods are absent or hormone levels suggest hypothalamic suppression.

Some clinics use BMI thresholds for certain treatments, usually because of anesthesia safety, procedural risk, or pregnancy risk rather than because pregnancy is impossible. These thresholds can feel distressing or stigmatizing. If you encounter a BMI policy, you have the right to ask for a clear explanation, alternatives, referral options, and a plan that supports your goals while prioritizing safety.

### **Weight and pregnancy health after conception**

Preconception weight matters not only for becoming pregnant but also for pregnancy health. Higher BMI is associated increased risk of complications such as gestational diabetes, hypertensive disorders of pregnancy, cesarean birth,

anesthesia complications, and miscarriage. The American Society for Reproductive Medicine notes that obesity is linked higher miscarriage rates and pregnancy complications including gestational diabetes.

Low BMI concerns, including nutritional deficiencies, inadequate gestational weight gain, and potential effects on fetal growth, depending on the individual situation. This is why includes assessment of nutrition, folic acid intake, or iron status ]] indicated, chronic disease management, medication review, and mental health support.

The goal is not to achieve a perfect number on a scale. The goal is to enter pregnancy as ly supported as possible. For some people, that may involve gradual weight loss; for s, weight restoration; for s, focusing on metabolic markers, regular meals, strength, sleep, or treatment of conditions such as diabetes, thyroid disease, PCOS, or eating disorders.

### **Healthy weight change before pregnancy: what support can look like**

If weight may , a supportive plan should be realistic, non-punitive, and ]]]y safe. Crash dieting, extreme exercise, unregulated supplements, or very low-calorie plans can worsen hormonal function and may be unsafe . Similarly, attempts to gain weight should focus on , adequate energy intake, and treatment of any underlying cause rather than simply increasing calories without context.

Helpful professional support may include:

A primary care clinician or obstetrician-gynecologist to review ]] history, , cycles, and risks.

A endocrinologist if has not occurred after an appropriate interval or if cycles are irregular.

An accredited planning, whether the goal is weight loss, weight gain, or metabolic optimization.

Mental heh support if eating, , stress, or ]] treatment is causing distress.

Screening and management for diabetes, thyroid disease, PCOS, sleep apnea, or other conditions when clinically indicated.

Even modest changes may improve ovulatory function or metabolic health for some people, but the appropriate target varies. If you are older than 35, have very

irregular cycles, have known conditions, or have been trying for some time, it may be better to seek [ ] while , rather than delaying care indefinitely to reach a specific BMI.

### **When to ask for fertility evaluation**

General guidance is to seek evaluation after 12 of regular unprotected intercourse if you are under 35, or after 6 if you are 35 or older. Earlier assessment is reasonable if you have absent or very irregular periods, known endometriosis, PCOS, loss, prior pelvic infection, chemotherapy exposure, significant medical illness, or a known -related factor.

If BMI is a concern, evaluation be comprehensive. A clinician may assess , ovarian reserve, tubal , uterine anatomy, semen parameters, metabolic heh, and endocrine conditions. This matters because weight can contribute to challenges, but it is not the only possible cause. A full assessment unnecessary delay and help tailor care.